

Flow Synthesis and Application of 2,2,5,5-Tetramethyltetrahydrofuran in Organic Reactions



Bernice M. Currie ^a, Jenny-Lee Panayides ^b, and Darren L. Riley ^{*a}

^a Department of Chemistry, University of Pretoria, Pretoria, 0028, South Africa;

^b Pharmaceutical Technologies, Chemicals cluster, CSIR, Pretoria, 0001, South Africa

Correspondence: Bernice.currie@tuks.co.za; Darren.riley@up.ac.za



ReactWise





1 Introduction

- Growing environmental and regulatory concerns surround the widespread use of petrochemical solvents (e.g. toluene), which are toxic and volatile yet remain essential in industry.^{1,2}
- TMTHF (TMO) is a safer alternative with similar properties and no peroxide formation, but application in organic chemistry has been limited.³
- Current TMTHF production relies on high cost and energy consumption and corrosive acids/zeolites, limiting sustainability and scalability.³
- Citric acid is a renewable, biodegradable, low-cost green catalyst.⁴
- Flow chemistry improves safety, efficiency, scalability, and sustainability.⁵

This work presents a simple, cost-effective, and scalable flow route to TMTHF using citric acid and water, and its thermal and photochemical stability in organic chemistry.

3 TMTHF synthesis



Flow conditions

- Conc. 0.75 M
- Flow rate: 2 ml min⁻¹
- Residence time: 20 minutes
- Temperature: 140 °C
- Run time: 15.25 hr

Results

- TMTHF is immiscible with water
- Yield: 125.7 grams (72%)
- Purity: 90%

Cost analysis:

- Space-time yield: 9.43 kg L⁻¹ h⁻¹
- Throughput: 8.24 g h⁻¹
- Price: 0.13 USD per gram

5 Claisen Rearrangement



Scheme 2. Claisen rearrangement in TMTHF under flow.

Table 2. Claisen rearrangement results using TMTHF.

Entry No	Conc. [M]	Flow rate (ml/min)	Tres (min)	Yield (%)
1	1.00	2.0	10	88.0
2	1.50	2.0	10	96.9
3	2.00	2.0	10	82.0
4	1.00	1.3	15	96.9
5	1.00	1.0	20	97.7
6	1.50	1.0	20	97.8
7	2.00	1.0	20	94.9
8 ^a	1.50	1.0	20	96.4

^a 50-gram reaction results

Throughput (g h ⁻¹)	16.72 g h ⁻¹
Space-time yield	7.23 kg h ⁻¹ L ⁻¹
Atom economy (%)	96.42 %

Claisen Rearrangement: Solvent Dielectric Constant Vs Yield





Chart 1: Effect of solvent dielectric constant on reaction performance.


6 Conclusion

- **TMTHF synthesis:** TMTHF was produced in flow on a multigram scale (125.7 g) using citric acid and water with a throughput of 8.24 g h⁻¹, and an estimated cost of 0.13 USD g⁻¹.
- **Photochemical application:** Nitroalkylation was optimized using ReactWise machine learning, identifying optimal conditions of 450 nm irradiation, 45 min residence time, 0.026 M concentration, and 2.93 equiv BrCCl₃, achieving 89.8% conversion, notably without the need for a photocatalyst.
- **Thermal application:** The Claisen rearrangement was optimized and scaled to 50 g, demonstrating solvent applicability with a yield of 96.4% and throughput of 6.72 g h⁻¹.
- **Solvent behavior:** Based on dielectric constant and reaction outcomes, TMTHF behaves more similarly to hydrocarbon solvents than traditional ether solvents, which may explain the improved performance observed in certain transformations.

2 Methodology



4 Photochemical Reaction



Scheme 1. Photochemical nitroalkylation of N-phenyl-tetrahydroisoquinoline in TMTHF (no photocatalyst).

Table 1. Nitroalkylation optimization results using ReactWise

Entry No	Type	λ (nm)	Tres (min)	Conc. [M]	BrCCl ₃ (Equiv.)	Conv. (%)
1	Initialize	365	14	0.038	1.76	59.15
2	Initialize	410	44	0.249	2.92	69.08
3	Initialize	450	28	0.161	0.58	75.84
4	Initialize	410	6	0.098	0.54	66.70
5	Initialize	365	39	0.211	1.09	52.96
6	Initialize	450	10	0.064	2.51	81.82
7	Initialize	410	35	0.029	2.15	62.55
8	Initialize	450	41	0.237	2.77	69.83
9	Optimize	450	45	0.026	2.93	89.82
10	Optimize	450	24	0.059	3.00	81.64
11	Optimize	450	34	0.026	3.12	86.69




Figure 1: 3D Surface Plot




Figure 2: Parameter importance plot




Figure 3: Categorical comparison plot at optimal model conditions

7 References

1. T. Welton, Solvents and sustainable chemistry, Proc. R. Soc. A. 471, 2183 (2015) 1-25. DOI: 10.1098/rspa.2015.0502.
2. F. G. Calvo-Flores, M. J. Montegudo-Arrebola, J. A. Dobado and J. Isac-Garcia, Green and Bio-Based Solvents, Top. Curr. Chem. 376, 18 (2018) 1-40. DOI: 10.1007/s41061-018-0191-6.
3. F. Byrne, B. Forier, G. Bossaert, C. Hoebets, T. J. Farmer, J. H. Clark and A. J. Hunt, 2,2,5,5-Tetramethyltetrahydrofuran (TMTHF): A non-polar, non-peroxide forming ether replacement for hazardous hydrocarbon solvents, Green Chem. 19 (2017) 3671-3678. DOI: 10.1039/C7GC01392B.
4. R. Ciriminna, F. Meneguzzo, R. Delisi and M. Pagliaro, Citric acid: Emerging applications of key biotechnology industrial product, Chem. Cent. J. 11, 22 (2017) 1-9. DOI:10.1186/s13065-017-0251-y.
5. S. G. Newman and K. F. Jensen, The role of flow in green chemistry and engineering, Green Chem. 15 (2013) 1456-1472. DOI: 10.1039/C3GC40374B.



Scan the QR code to access the full published article on the synthesis of TMTHF.